P510/1 **PHYSICS** Paper 1 July/Aug. 2018 21/2 hrs

INTERNAL MOCK EXAMINATINS – 2018

Uganda Advanced Certificate of Education

Physics Paper 1

2 hours 30 minutes

INSTRUCTIONS TO CANDIDATES:

Attempt five questions, including at least one but not more than two questions from each of the sections A, B and C. Any additional question(s) answered will **not** be marked.

Non-programmable scientific calculators may be used.

Assume where necessary:

 $9.81 \, \text{ms}^{-2}$ Acceleration due to gravity, g

 $1.6 \times 10^{-19} C$ Electronic charge, e

 $9.11 \times 10^{-31} kg$ Electron mass

 $5.97 \times 10^{24} kg$ Mass of the earth

 $6.6 \times 10^{-34} Js$ Plank's constant. h

 $5.67 \times 10^{-8} WM^{-2} K^{-4}$ Stefan's – Boltzmann's constant σ

 $6.4 \times 10^{6} m$ Radius of the earth

 $7 \times 10^{8} m$ Radius of the sun

 $1.5 \times 10^{11} m$ Radius of earth's orbit about the sun

 $3.0 \times 10^8 \text{ms}^{-1}$ Speed of light in a vacuum, C

390Wm⁻¹K⁻¹ Thermal conductivity of copper

 $210Wm^{-1}K^{-1}$ Thermal conductivity of aluminium

4200Jkg⁻¹K⁻¹ Specific heat capacity of water

 $6.67 \times 10^{-11} \, \text{Nm}^2 \text{kg}^{-2}$ Universal gravitational constant, G

 $6.02 \times 10^{23} mol^{-1}$ Avogadro's number, N_A

 $7.0 \times 10^{-2} \text{Nm}^{-1}$ Surface tension of water

Density of water $8.31.Imol^{-1}K^{-1}$ Gas constant, R

 $1.8 \times 10^{11} \, \text{Ckg}^{-1}$ Charge to mass ratio, e/m

The constant, $\frac{1}{4\pi\varepsilon_0}$ $9.0 \times 10^{9} \text{F}^{-1} \text{m}$ =

400JKg⁻¹K⁻¹. Specific heat capacity of copper

 $3.3 \times 10^{5} J K g^{-1}$. Specific latent heat of fusion of ice

 $9.56 \times 10^{4} Cmol^{-1}$ Faraday constant, F

TurnOver

1000Kgm⁻³

SECTION A

- 1. (a) (i) State Archimedes principle. (01 mark)
 - (ii) Describe an experiment to determine the relative density of an irregular solid which floats in water. (03 marks)
 - (b) A solid weighs 237.5g in air and 12.5g when totally immersed in a fluid of density $900Kgm^{-3}$. Calculate the density of the liquid in which the solid would float with one fifth of its volume exposed above the liquid surface.

(06 marks)

- (c) (i) What is meant by viscosity. (01 mark)
 - (ii) Explain the effect of temperature on the viscosity of a liquid. (03 marks)
- (d) (i) State the work energy theorem. (01 mark)
 - (ii) A bullet of mass 100g moving horizontally with a speed of $420ms^{-1}$ strikes the block of mass 2000g at rest on a smooth table and becomes embedded in it. Find the kinetic energy lost if they move together.

 (05 marks)
- 2. (a) (i) Define centre of gravity. (01 mark)
 - (ii) Describe an experiment to find the centre of gravity of a regular piece of card board. (03 marks)
 - (b) Explain using the molecular theory the laws of solid friction. (07 marks)
 - (c) (i) Define surface tension. (01 mark)
 - (ii) Explain the origin of surface tension. (03 marks)
 - (d) Explain why rain drops hit the ground with less force than they should. (05 marks)
- 3. (a) State Newton's laws of motion. (03 marks)
 - (b) A body X of mass m_1 moves with velocity u_1 and collides head on elastically with another body, Y of mass m_2 which is at rest. If the velocities of X and Y are v_1 and v_2 respectively and given that $A = m_1/m_2$ show that;
 - (i) $\frac{u_1}{v_1} = \frac{A+1}{A-1}$ (04 marks)

	$\frac{v_2}{v_1} = \frac{2A}{A-1}$	(03 marks)
` /	v_1 $A-1$	` '

- (c) Describe an experiment to determine the acceleration due to gravity using a spiral spring of known force constant. (05 marks)
- (d) Explain the following:
 - (i) A mass attached to a string rotating at a constant speed in a horizontal circle will fly off at a tangent if the string breaks. (02 marks)
 - (ii) A cosmonaut in a satellite which is in a free circular orbit around the earth experiences a sensation of weightlessness even though there is influence of gravitational field of the earth. (03 marks)
- 4. (a) (i) What is meant by simple harmonic motion? (01 mark)
 - (ii) State **four** characteristics of simple harmonic motion. (02 marks)
 - (b) A mass, m is suspended from a rigid support by a string of length, X. The mass is pulled a side so that the string makes an angle, θ with the vertical and then released. Show that the mass executes simple harmonic motion with a period $T = 2\pi \sqrt{\frac{x}{g}}$. (05 marks)
 - (c) A horizontal spring of force constant $300Nm^{-1}$ fixed at one end has a mass of 3kg attached to the free end and resting on a smooth horizontal surface. The mass is pulled through a distance of 5.0cm and released. Calculate;
 - (i) angular speed, (02 marks)
 - (ii) maximum velocity attained by the vibrating body. (02 marks)
 - (iii) acceleration when the body is half way towards the centre from its initial position. (02 marks)
 - (d) (i) What is meant by a couple in mechanics? (01 mark)
 - (ii) State the conditions for equilibrium of a system of coplanar forces. (02 marks)
 - (e) Explain why a person standing near a railway line is sucked towards the railway line when a fast moving train passes. (03 marks)

SECTION R

			SECTION D				
5.	(a)	(i)	Define thermal conductivity.		(01 mark)		
		(ii)	Explain the mechanism of heat transfer in n	netals.	(03 marks)		
	(b)	by a	A double glazed window has two glass sheets of thickness $5.0mm$, separated by a layer of air of thickness $1.2mm$. If the two inner air – glass surfaces have steady temperatures of $25^{\circ}C$ and $.5^{\circ}C$ respectively, find the:				
		(i)	temperature of the outer – glass surfaces.	(03 n	narks)		
		(ii)	amount of heat that flows across an area hours.	of the wind	ow of $3m^2$ in 3 (03 marks)		
		(Conductivity of glass = $0.72Wm^{-1}K^{-1}$ and that of air = $0.025Wm^{-1}K^{-1}$					
	(c)	(i)	What is a perfectly black body?	(01 mark)			
		(ii)	The energy intensity received by a sphoton $1.5 \times 10^3 Wm^{-2}$, The star is of radius 7.0	-			
	1.4 star.	$1.4 \times 10^8 Km$ from the planet. Calculate the surface temperature of star. (04 marks)					
	(d)	Expl	lain the green house effect and how it is relate	ed to global w	varming. (05 marks)		
6.	(a)	Defi	ne specific heat capacity of a substance.		(01 mark)		
	(b)	(i)	Describe an electrical method for the determination of a metal.	nination of th	ne specific heat (06 marks)		
		(ii)	State the assumptions made in the above ex	periment.	(02 marks)		
	(c)	340, and	Steam at $100^{\circ}C$ is passed into a copper calorimeter of mass $150g$ containing $340g$ of water at $15^{\circ}C$. This is done until the temperature of the calorimeter and its content is $71^{\circ}C$. If the mass of the calorimeter and its contents is found to be $525g$ calculate the specific latent heat of vaporization of water. (06 marks)				
	(d)	(i)	State the assumptions made in the derivatio	n of the expr	· · · · · · · · · · · · · · · · · · ·		
			$P = \frac{1}{3} fC^{-2}$ for the pressure of an ideal gas	s. (02 n	narks)		
		(ii)	Use the expression in (d) (i) above to deduce pressures.	e Dalton's la	w of partial (03 marks)		

- 7. (a) Define a thermometric property and give **two** examples. (02 marks)
 - (b) The resistance, R_{θ} of platinum varies with temperature $\theta^{o}C$ as measured by a constant volume gas thermometer according to the equation:

$$R_{\theta} = 50.0 + 0.17\theta + 3.0 \times 10^{-4}\theta^{2}$$

- (i) Calculate the temperature on the platinum scale corresponding to $60^{\circ}C$ on the gas scale. (06 marks)
- (ii) Account for the difference between the two values and the temperature at which they agree. (02 marks)
- (c) Use the kinetic theory of matter to explain the following observations:
 - (i) Saturated vapour pressure of a liquid increases with temperature. (03 marks)
 - (ii) Saturated vapour is not affected by a decrease in volume at constant temperature. (03 marks)
- (d) An ideal gas of volume $100cm^3$ at s.t.p expands a diabatically until its pressure drops to a quarter its original value. Find the new volume and temperature if the ratio of the principal specific heat capacities is 1.4 (04 marks)

SECTION C

- 8. (a) (i) What is meant by the terms: radio active, decay, half life and decay constant? (03 marks)
 - (ii) Show that the half—life $t_{\frac{1}{2}}$ of a radio isotope is given by $t_{\frac{1}{2}} = \frac{693}{\lambda}$ where λ is the decay constant. (Assume the decay law $N = Noe^{-\lambda t}$).
 - (b) With the aid of a diagram describe the structure and action of a diffusion cloud chamber. (05 marks)
 - (c) The radio isotope ${}_{38}^{90}X$ decays by emission of β particles. The half life of the isotope is 28.8 year. Determine the activity of 1g of the isotope.(05 marks)
 - (d) (i) What are cathode rays? (01 mark)
 - (ii) An electron accelerated by a p. d of 1000V passes through a uniform electric field intensity crossed with a uniform magnetic field of flux density 0.3T. If the electron emerges undeflected, calculate the electric field intensity. (03 marks)

- 9. (a) (i) Define space charge as applied to thermionic diodes. (01 mark)
 - (ii) Draw anode current anode voltage curves of a thermionic diode for two different filament currents and explain their main features.

(06 marks)

- (b) Derive the expression for the amplification factor μ in terms of anode resistance, R_a and mutual conductance, g_m for a triode valve. (03 marks)
- (c) A triode with mutual conductance $3mAV^{-1}$ and anode resistance of $10k\Omega$ is connected to a load resistance of $20k\Omega$. Calculate the amplitude of the output signal if the amplitude of the input signal is 30mV. (04 marks)
- (d) (i) What is a photon? (01 mark)
 - (ii) Explain using the quantum theory, the experimental observations on photo electric effect. (05 marks)
- 10. (a) A beam of α particles is directed normally to a thin gold foil. Explain why
- (i) most of the alpha particles past straight through the foil, (02 marks)
 - (ii) few alpha particles are deflected through angles more than 90° . (02 marks)
 - (b) Calculate the least distance of approach of a 4.0*MeV* alpha particles to the nucleus of a gold atom. (Atomic number of gold = 79). (04 marks)
 - (c) Explain, using suitable sketch graphs, how X ray spectra in an X ray tube are produced. (06 marks)
 - (d) A beam of X rays of wave length $9 \times 10^{-11} m$ is incident on a sodium chloride crystal of interplanar separation $9.0 \times 10^{-10} m$. Calculate the first order diffraction angle. (03 marks)
 - (e) (i) Distinguish between nuclear fussion and nuclear fission. (02 marks)
 - (ii) State the conditions necessary for each of the nuclear reactions in (e) (i) to occur. (01 mark)

END