UACE - UNNASE 2019.

Write on both sides of the paper | Question | Do not write in this margin

UNNASE 2019 BIOLOGY PAPER II GUIDE;

ION

i a) Comparison of the effect of increasing light intensity on the net rate of oxygen release by the 2 plant types.

Similarities;

Net rate of oxygen release for both plant types attain a compensation point; ✓

The net rate of oxygen release for both plants types increases with increasing light intensity or photon irradiance up to 80 μmol quanta $m^{-2}s^{-1}$; ✓

Both plant types attain the same net rate of oxygen release at a light intensity of 100 μmol quanta $m^{-2}s^{-1}$; ✓ any 3 (03)

A maximum net rate of oxygen release was attained by both plant types.

Differences;-

Shade plant	Sun plant.
- Compensation point attained at a lower light intensity	- Compensation point attained at a higher light intensity; ✓
- Maximum net rate of oxygen release lower.	- Maximum net rate of oxygen release higher; ✓
- Maximum net rate of oxygen release attained at lower light intensity	- Maximum net rate of oxygen release attained at a higher light intensity; ✓
- Net rate of oxygen release constant from light intensity of = 80 μmol quanta $m^{-2}s^{-1}$.	- Net rate of oxygen release increases rapidly and then gradually from a light intensity of = 80 μmol quanta $m^{-2}s^{-1}$; ✓ (05)
- Net rate of oxygen release higher at low light intensity	- Net rate of oxygen release lower at low light intensities; ✓ any 5
- Net rate of oxygen release lower at high light intensity	- Net rate of oxygen release is higher at high light intensity;

ii) Effect of increasing light intensity on the Sun plant.

- Slight increase / Big increase in overall net rate of O_2 release;
from a = 80 μmol quanta $m^{-2}s^{-1}$; net rate of oxygen release increased gradually but still below compensation point; implying oxygen was consumed since the light intensity was a strong limiting factor; for photosynthesis hence causing more photolysis but oxygen used for

(1)

08

respiration was more than the oxygen produced; from the slight photolysis of water in the light dependent stage of photosynthesis.
At a light intensity of ≈ 80 μmol quanta $m^{-2} s^{-1}$, a compensation point was attained; implying net rate of oxygen release was equal to net rate of oxygen consumption; implying that the photolysis rate had increased to make net rate of oxygen release and consumption to be equal.; max (07)
from approximately 100 to 400 μmol quanta $m^{-2} s^{-1}$; net rate of oxygen release increased rapidly above the compensation point; implying that light intensity was a pronounced limiting factor still; and so its increase rapidly increased the rate of photosynthesis or water photolysis hence producing more oxygen than was what is consumed in respiration;
The net rate of oxygen release then increased gradually with further increase in light intensity; from ≈ 400 to 1000 μmol quanta $m^{-2} s^{-1}$, implying that light intensity was no longer the major limiting factor of photosynthesis; and instead other factors such as temperature, water, mineral salts, carbondioxide concentration and chlorophyll concentration; were limiting the photolysis of water in photosynthesis.

iii) Suggestion and explanation for the difference in effect of increasing light intensity on the two plants;
At low light intensity, the shade plant had a higher rate of oxygen release; higher maximum rate of photosynthesis; and attained its compensation point; because shade plants are photosynthetically more efficient at such low light intensities. This is because the shade plants possess a lower light saturation point; and are more adapted to harvest and harness low light intensity; by possessing broader leaves; and more chlorophyll since their chloroplasts possess more grana.;
Sun plants on the other hand had a lower net rate of oxygen release at low light intensities, but to their maximum net rate of oxygen release, and compensation point were attained at higher light intensities because they are photosynthetically more efficient at higher light intensities;

This is because they are well adapted to harvest and harness high light intensity due to their narrower thick leaves or lamina, many layers of palisade mesophyll cells, and possession of chloroplasts which take turns in bright light and shelter in others' shade as they use the absorbed light to prevent Chlorophyll destruction; ✓ (08)

The sun plants also had a higher maximum rate of net oxygen release; and a greater effect of increasing light intensity on their net rate of oxygen release; because sun plants have a higher light saturation point than the shade plants; ✓

b(i) Explanation for effect of increasing CO_2 concentration and temperature on net CO_2 assimilation by barley.

Increasing Carbondioxide concentration;

The net CO_2 assimilation rate of barley increased with increase in CO_2 level or concentration because barley is a C_3 plant and increase in CO_2 level reduces the inhibitory rate of oxygen on RUBISCO; hence reducing photorespiration which is wasteful; since it produces less phosphoglycerate, utilizes ATP and $NADPH_2$ and reduces CO_2 combination with RUBISCO. The increase in CO_2 level also increases chances of collision with RUBISCO; which increases the rate of photosynthesis and hence net rate of CO_2 assimilation. (01½)

Increasing temperature;

Increase in temperature from 5 to 15°C caused a gradual increase in net rate of CO_2 assimilation for barley plants grown in both ambient CO_2 and high CO_2 levels; due to activation of the photosynthetic enzymes; which activate photosynthesis;

Increase in temperature from 15 to 30°C caused a gradual ma decrease in net CO_2 assimilation rate for barley plants growing in ambient CO_2; due to the CO_2 levels limiting the assimilation rate; which increases photorespiration; since O_2 molecules are lighter, hence and more are formed collide more with temperature increase and CO_2 limitation; (03½)

The same temperature increase caused a rapid increase in net CO_2 assimilation for Barley plants grown in high CO_2 environments; (3)

due to more activation of the enzymes with CO_2 abundance and due to low or no photorespiration since the CO_2 levels are high hence competitively out competing O_2 for the active sites of RUBISCO (Ribulose bisphosphate carboxylase oxygenase).

iii) Explanation for the difference in Net CO_2 assimilation rate for maize and Barley at ambient CO_2 levels.

The Net CO_2 assimilation rate is higher for Barley at low temperatures because barley is a C_3 plant whose photosynthetic enzymes are more efficient at lower temperatures, unlike maize which is a C_4 plant whose photosynthetic enzymes are inefficient at such low temperatures. 05

Increase in temperature from 20°C to 30°C caused a higher net CO_2 assimilation rate in maize due to increased efficiency of maize's photosynthetic enzymes and decreased efficiency for Barley's photosynthetic enzymes and due to increased photorespiration in Barley plants with such temperatures.

iii) From figure 2 why C_4 plants tend to be found in hotter more arid regions than C_3 plants.

From figure 2 (two) below 15°C, net rate of CO_2 assimilation is very low for maize which is a C_4 plant due to the low efficiency of their photosynthetic enzymes at such temperatures hence low productivity.

As the temperatures increased above 15°C which is characteristic of hotter more arid regions, the net CO_2 assimilation rate increased rapidly due to increased photosynthetic enzymes' activation and hence increased photosynthesis, primary productivity and survival. 07 max

From figure 2, C_3 plants' Net CO_2 fixation or assimilation rate was higher in temperatures below 15°C due to high efficiency of the enzymes and less photorespiration and the rate of Net CO_2 assimilation decreased with increase in temperatures beyond 15°C due to reduced enzymes' activation or denaturation and more photorespiration in such hotter conditions; (4)

2(a) Description of the physiological behaviour of rods during:

i) Darkness:

The rods are not stimulated; and the sodium pump in its inner segment membrane of the outer segment actively pumps out sodium ions; Since the outer membrane is leaky; some sodium ions diffuse back in to the rod; down their concentration gradient there by reducing the negative charge inside the rod which maintains the resting potential;

This results in maximum release of the neurotransmitter substance usually glutamate; at the terminal end of the rod with which it synapses with the bipolar neuron. This maintains a resting potential in the bipolar neuron; and no impulses are fired via the ganglion cells of the optic nerve; (05)

ii) Low light intensity;

The rods are stimulated; resulting in bleaching or isomerisation; of cis retinene in rhodopsin to trans-retinene; which causes photo-decomposition; of rhodopsin into opsin/scotopsin and retinene; in a process of bleaching;

This leads to development of the generator potential; upto a threshold; there by reducing the permeability of the outer membrane of the outer segment towards sodium ions; with continued active pumping of sodium ions out of the outer segment; by the sodium pump; with in the inner membrane of the segment of the rod, hyperpolarisation of the rod results; (08)

This decreases the rate at which the neurotransmitter substance of glutamate is released at the terminal rod; leading to hyperpolarisation of the bipolar neurons; hence propagation of an action potential; along the optic nerve by the ganglion cell;

iii) Excessively high light intensity;

The rhodopsin in the rods is over bleached; above by the rate of its photodecomposition into scotopsin/opsin and retinene is higher than the rate of resynthesis or regeneration.; (5)

This renders the amount of rhodopsin insufficient and cannot be temporarily used to focus objects in dim light as the individual moves to low light intensity regime. (02)

b) Significance of the compactness of cones at the fovea.

By being closely packed together, the cones increase visual acuity or precision; in that the images of two close objects can fall on two different cones; seperated by at least one cone; which results into interpretation of images as seperate units; of images by the brain; (02)

20

3(a) Explanation for determination of:

i) ABO blood groups;

This is determined by an autosomal gene; whose locus is represented by a symbol I standing for isohaemagglutinogen;

The gene has more than three alleles or has multiple alleles; A, B and O; whose by alleles A and B are codominant, while O is recessive to both A and B;

When an individual inherits any of the following allelic combinations via fertilization, he or she tends to possess any of the (02)
the 4 blood groups A, B, AB and O;

Genotype $I^A I^A$ or $I^A I^O$ determine formation of agglutinin 'b' (or antibody) in plasma and antigen/agglutinogen A; on the erythrocyte and the individual has blood group A;

Inheritance of genotype BB or BO leads to production of agglutinin 'a' in plasma and antigen B on erythrocytes hence possession of blood group B;

Genotype $I^A I^B$ leads to production of both alleles A and B; on erythrocytes without agglutinins/antibodies; in plasma which render the individual to have blood group AB;

Genotypes $I^O I^O$ codes for production of agglutinins a and b; with our antigens; enabling the individual to be blood group O;

(b) (ii) Sex in man;

Sex in man is determined by Sex chromosomes; X and Y; carried in gametes. During fertilization, an embryo may either inherit

XX genotype ; of fusing sperm contain the X- Sex Chromosome; for
XY genotype of the fusing sperm contain the Y- Sex Chromosome;
The XX genotype determines the female homogametic sex; while
the XY genotype determines the male heterogametic sex.;
Since all embryos possess an X Chromosome which carries the testicular feminisation gene; for production of a protein in form of testosterone receptor, all of them have this receptor. Because the would be male embryos have a Y Chromosome with a testes determining gene; for production of a protein called H-Y antigen / Testes determining factor (TDF); on the body cells of a developing male. The non differentiated gonads / genital ridges differentiate into testes / Semeniferous tubules + interstitial cells; max (07)
- Kiththis the H-Y antigen / TDF, the non differentiated gonads differentiate into ovary of a female embryo; the developing testes of the male embryo secrete testosterone; into the blood stream which binds on the testosterone receptor to form the testosterone - receptor complex; which moves from the cytoplasm of the target cell via the nuclear pore into the nucleus and it activates the genes which code for development of the tissues which give rise to the male reproductive system; the tissues of a developing female embryo are not activated due to lack of testosterone hence a female reproductive system develops.;

b) Evolution of Charles Darwin's finches;
Originally, there was one predominant; mainland ground finch variety with short straight beaks; for feed and crushing. and insects
There were few other variants of finches regarding beak shape and size; like the cactus finches with long slightly curved beak and a split tongue to feed on nectar, Insectivorous finch with parrot like beak to feed on seeds and insects, vegetarian finch with curved parrot like beak to feed on fruits, buds and soft fruits, warbler finch with slender beak to feed on flying insects and the wood pecker finch with large straight beak to use cactus spine to poke insects out of holes in wood; any other like.

The high population of the finches on the main land resulted in stiff competition for the limited resources. This caused migration of some finches to the vacant Galapagos Islands whose niches were different especially regarding the predominant food types.
The finch varieties whose beak shapes and sizes best suited exploiting the different unique food types on the different specific Islands survived more, and reproduced more; while the less max suited were selected against hence survived less.
The adaptive features in the off springs got concentrated; aged with isolation of land masses, gene flow between finch varieties unique to the Islands was prevented; and over generations of genetic dissimilarity; different more adapted finch species per Island were developed or arose. 20

4 a) How Stratosphere is destroyed:
A Stratosphere is the upper layer of the atmosphere; 15-40 kms above the earth surface; with the Ozone layer; which absorbs short wave length radiations; like ultraviolet rays, x-rays and gamma rays; from the sun and prevent them from reaching the earth surface in large amount.
Ozone is destroyed by fluorine; and chlorine radicals; from the breakdown of chlorofluorocarbons (CFCs); using ultraviolet radiation from the sun. 13
The CFCs; are used as aerosol sprays; or coolants in refrigerators; while chloromethane; from rotting vegetation and bush burning; also react with Ozone in the Ozone layer.
When the fluorine and chlorine radicals react with ozone in using energy from ultraviolet radiations of sunlight, the Ozone is broken down into molecular oxygen at a rate faster than at which it is reformed.
Bromine; also gives off radicals which react and break down ozone. It is mainly obtained from methyl bromide used in fungicides; Nitrogen monoxide; from the reaction of dinitrogen oxide with atomic oxygen; using energy from U.V radiations also react with and

Write on both sides of the paper

Question

Do not write in this margin

destroy the Ozone layer; ✗

The high flying aircrafts release icy particles in the Stratosphere leading to depletion of the ozone layer; ✗

b) Indicators to support the destruction of the stratosphere:

The large amounts of short wave length radiations from the Sun reaching the earth surface; ✓

The more frequent change in the nucleic acids or mutations which result into genetic diseases and abnormalities eg skin cancers.

Crop damage which is characteristic of absorption of short wave length radiations is also common and is an indicator of ozone layer destruction. ✓ max 03

Regular destruction of aquatic species of organisms mainly the fish larvae and plankton; is indicative of increased short wave length radiation from the Sun which is evidence of stratosphere destruction. ✓

Reduced carbondioxide absorption by algae resulting in increased global warming is evidence of algal destruction by short wave length radiations from the sun hence indicative of stratosphere destruction. ✓

c) An indicator species.

This is a species of organisms which are highly sensitive to a certain environmental factor, shown by their presence, absence or relative numbers, reflects the level or absence of that factor in the environment; ✗ like the Lichens are indicator species of the (consider any appropriate eg. Concentration of Sulphurdioxide in the atmosphere because Lichens are highly poisoned and killed by Sulphurdioxide, their large population in an area reflects low levels or absence of SO_2; in the area and its low population reflects high levels of SO_2 in the area; (Consider any approximate other example). ✓ max 04

20

b) Effect of light on morphology of plants: (NB red light = Sunlight)
Red light stimulates leaf expansion in dicots while far-red light inhibits it;

Red light stimulates unrolling of leaves of grasses while far-red light inhibits it.

Red light stimulates chloroplast development; while far-red light inhibits it hence causes chlorosis

Red light inhibits internode growth; while far-red light stimulates it

Darkness or total lack of light causes elongation of internodes; while light inhibits it.

Darkness causes development of embryonic leaves; while light stimulates growth of adult leaves.

Far-red light stimulates lateral roots growth; while red light inhibits lateral roots growth.

Red light stimulates greening of all plant parts while far-red light inhibits it.

Red light also stimulates unhooking of plumule in dicots while far-red light inhibits it.

Blue light stimulates increase in stem diameter of plants, while other light qualities have no significant effect.

Blue light also causes increase in general fresh weight of the plant and other light wave lengths have no effect.

Exposure of a plant to blue light quality increases stomatal frequency which thus changes morphology.

10

20